

Scopus

by Toni Malvin

Submission date: 25-Feb-2023 08:10AM (UTC-0600)

Submission ID: 2022739079

File name: 169-174.pdf (733.79K)

Word count: 5704

Character count: 28960



Effects of Dietary Catechin *Uncaria gambir* Extract on Growth Performance, Carcass Characteristics, Plasma Lipids, Antioxidant Activity and Nutrient Digestibility in Broiler Chickens

Ramaiyulis^{1*}, Mairizal², Salvia¹, Nelzi Fati¹ and Toni Malvin¹

¹Animal Production Technology Study Program, Agricultural Polytechnic Payakumbuh, Lima Puluh Kota 26271, Indonesia

²Department of Poultry Nutrition, Faculty of Animal Science, Universitas Jambi, Jambi, Indonesia

*Corresponding author: ramaiyulis@gmail.com

Article History: 22-613

Received: 14-May-22

Revised: 18-Jun-22

Accepted: 22-Jun-22

ABSTRACT

This study aimed to determine the effect of adding catechin gambir extract (*Uncaria gambir* Roxb) (CGE) on growth performance, carcass characteristics, blood plasma constituents, antioxidant activity, and nutrient digestibility in broiler chickens. 160 chicks were allocated to receive four treatment diets adding CGE was 0, 0.1, 0.2, and 0.3g/kg in basal diets with four replications. Bodyweight and feed intake were recorded to calculate growth performance. Digestibility measurements were carried out at the age of 31-40 days, and at 40 days, blood samples were taken to determine the total protein levels of cholesterol, low-density lipoprotein (LDL) cholesterol, high-density lipoprotein (HDL), and Malondialdehyde (MDA). Then these chickens are slaughtered to determine the percentage of viscera and muscles. Adding CGE to basal diets linearly ($P<0.05$) increased body weight gain and improved the feed conversion ratio of broiler grower, finisher, and the overall age of 1-40 days. The effect of GCE on carcass characteristics significantly ($P<0.05$) increased the percentage of thigh and drumstick and decreased abdominal fat. CGE linearly reduced total cholesterol and LDL while increased HDL and total protein in blood plasma. The MDA concentration linearly decreased in serum and broiler meat aged 40 days. In addition, CGE increased ($P<0.05$) the digestibility of crude protein and decreased fat digestibility in the diet. Adding 0.3g/kg CGE to the basal diet can be a feed additive to produce better performance and broiler meat.

Key words: Antioxidants, Broiler, Catechin, Digestibility, Malondialdehyde, *Uncaria gambir*

INTRODUCTION

The broiler industry plays an essential role in supplying meat as a source of animal protein for humans. Therefore, the meat produced must be healthy and free from harmful substances such as antibiotic residues and hormones, which concern using antibiotic growth promoters (AGP) (Lillehoj et al. 2018). Since the ban on AGP in many countries, medicinal plants as antibiotics and natural growth promoters in poultry diets have gained much attention (Solangi et al. 2020; El-Hack et al. 2020). In addition, medicinal plants can be used as alternatives to AGP as they exhibit antimicrobial properties and thus can become an integral part of poultry nutrition (Daramola 2019).

Flavonoids are compounds widely recommended as a substitute for AGP due to their ability as antibacterial, antioxidant, anti-inflammatory, and hepatoprotective. Therefore, they provide better broiler performance in the form of increased body weight gain and feed efficiency (Giannenas

et al. 2018) and they also improve meat quality (Liu et al. 2020) and reduce mortality (Prihambodo et al. 2021).

Catechin is a plant bioactive flavonoid compound that can be used as a broiler feed additive. There have been no reports of pure catechins in broiler diets. The inclusion of green tea (13.93% of catechins) as a feed additive in broilers can increase feed conversion ratio (FCR), decrease abdominal fat, and increase antioxidant status (El-Hack et al. 2020). Grape pomace (0.58mg/g of catechin) in a broiler diet can improve gut morphology and modify the cecal bacterial community and biochemical blood profiles without adverse effects on growth performance and meat quality (Erinle et al. 2022). Grapeseed (0.1g/g of catechin) is recommended as a synthetic antioxidant in broiler diets (Gungor et al. 2021).

The gambir is an extraction product of the gambir plant (*Uncaria gambir* Roxb) from the *Rubiacae* family, a potential catechin producer from Indonesia. Gambir contains many monomeric flavonoids, almost totally

Cite This Article as: Ramaiyulis, Mairizal, Salvia, Fati N and Malvin T, 2023. Effects of dietary catechin *Uncaria gambir* extract on growth performance, carcass characteristics, plasma lipids, antioxidant activity, and nutrient digestibility in broiler chickens. International Journal of Veterinary Science 12(2): 169-174. <https://doi.org/10.47278/journal.ijvs/2022.177>

catechins, and a trace of epicatechin (Widiyarti et al. 2020). Gambir catechin purification has been successfully carried out (Yeni et al. 2014; Ferdinal 2014) through 2-stage extraction. Gambir catechins are white to yellowish crystals and have a sweet taste (Ferdinal 2014). Therefore, we hypothesized that gambir catechins as a feed additive could cause positive effects on broiler nutrition. It is the first study to examine the effect of gambir catechin on growth performance, carcass characteristics, plasma lipids, antioxidant activity, and nutrient digestibility as an exploratory step for its use as AGP in broiler diets.

MATERIALS AND METHODS

Extraction and Purification of Gambir Catechins

Gambir was obtained from the gambir industry in Harau, Lima Puluh Kota District, West Sumatra, Indonesia. The catechin purification process was carried out by 2-stage extraction following the method suggested by Yeni et al. (2014) and Ferdinal (2014). In the first stage, gambir flour mashed by using a grinder machine with a 500-micron sieve, was macerated with ethanol for 3 days, then filtered, and the ethanol was evaporated using a rotary evaporator (Heidolph Laborota 4000) 280rpm rotation speed, temperature 50°C and pressure 460mbar, so that ethanol extract was produced. The second stage was extracted by adding 50% (w/v) distilled water by heating at 70°C with stirring for 20min. It was then filtered with a cotton cloth and then dried with a tunnel dryer at a temperature of 40°C to obtain catechin gambir extract (CGE), which was used as a feed additive.

Measurement of CGE Polyphenol Content

Measurement of CGE polyphenol content was carried out using the following method, and the measurement results are shown in Table 1.

The total phenolic content was determined by spectrophotometric methods (Chun et al. 2003). First, 1g of extract was added with 9mL of distilled water in a 25mL volumetric flask. Then 1mL of Folin-ciocalteu phenol reagent was added and allowed to stand for 5min. Then, 10mL of 7% Na₂CO₃ solution was added. Then the solution was diluted in a 25mL volumetric flask with distilled water and allowed to stand for 90min, and then the absorbance was measured at 750nm with a UV-Visible spectrophotometer (Shimadzu UV-1700) with the gallic acid standard.

The catechin content was determined following the method reported (Kassim et al. 2011) using HPLC. Samples were analyzed on a Shimadzu ADVP using a Chromolith SemiPrep RP-18 column (100-10 mm, Merck) at a flow rate of 0.5mL/min and detected at 280 nm using a UV detector. The chromatogram of a standard mixture of (+)-catechin hydrate, (-)-epicatechin, (-)-epicatechin gallate, (-)-galocatechin, and (-)-epigallocatechin was obtained by the gradient elution as described by Kassim et al. (2011). Two different mobile phases were prepared accordingly: Solvent A, acetonitrile/water (50:50 v/v) with 0.1% acetic acid; solvent B, acetonitrile/water (5:95 v/v) with 0.1% acetic acid. All mobile phases were filtered through PTFE Millipore (Whatman) 0.45µm filter paper and degassed with a sonicator for 10 minutes.

Total Condensed tannin contents were calculated following a modified method (Kassim et al. 2011). 100mg of catechin extract was dissolved in 10mL of distilled water, then 2mL of 5M HCl and 2mL of formaldehyde (Merck) 37% were added. The mixture was heated under reflux for 60min, then filtered. The reddish precipitate was washed with 10mL of hot water five times. The precipitate was then dried in a silica gel desiccator and weighed. Yield is expressed as a percentage of the weight of the starting material.

The antioxidant activity test was performed using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging method (Yeni et al. 2014). First, the 4mL extract solution was added with 1mL of 0.001M DPPH solution, then shaken until homogeneous and allowed to stand for 30 minutes. Ethanol was used as a control. Next, the absorbance of the solution was measured with a UV-Visible spectrophotometer (Shimadzu UV-1700) at a wavelength (λ) of 517nm. Finally, the control absorbance value calculated the inhibition percentage (IC50).

Chicken and Treatment Diet

The experimental procedure was approved by the Center for Research and Community Service No. 195/PL25/PL.00.02/2020, following Government Regulation of the Republic of Indonesia No. 95 of 2012 concerning veterinary public health and animal welfare.

A total of 160 DOC broiler strains of Loughmann MB 202 males were reared in an open house cage. Chicks were allocated to 16 cage boxes containing ten birds to receive four treatment diets with four replicates for 40 days.

The diet used was commercially produced by Japfa Comfeed, Indonesia with the nutritional composition of the diet shown in Table 2. The treatment added CGE in basal diets of commercial diets in the starter and finisher phase of broiler, while the pre-starter phase was not added. The addition of CGE in the diet had 4 levels, namely 0, 0.1, 0.2 and 0.3g/kg. CGE flour was added to the floured diet and stirred with a horizontal type mixer (1W-200, Indonesia) until homogeneous. Diet was provided in flour, and drinking water was provided ad libitum.

Bodyweight Measurement

For chickens aged 20 and 40 days, body weight and feed intake were measured after being fasted for 12 hours to determine average daily gain (ADG), average daily feed intake (FI), and feed conversion (FC) = FI/ADG.

Blood and Organ Sampling

At broiler aged 40 days, blood samples were taken from 8 cages randomly selected with two animals per cage. Blood samples were taken from the cutanea ulnar vein as much as 5mL. Blood sample preparation was carried out to obtain protein-free blood filtrate to determine serum glucose levels and total protein, cholesterol, LDL, HDL, and MDA levels using spectro-photometric techniques.

The Elitech group (2012) method calculated the total cholesterol and HDL of broiler blood serum. First, up to 10µL of broiler blood serum was pipetted and then poured into the test tube and 1000µL cholesterol reagent was added and mixed, then incubated for 10 minutes. Afterward, it was calculated by a photometer. Next, broiler blood serum was pipetted up to 250µL, the reagent of HDL up to 500µL

was added and centrifuged for 10 minutes at the speed of 2500rpm. Then, the supernatant was pipetted 100µL and added to the cholesterol reagent 1000µL, mixed and incubated for 10 minutes, then it was calculated by a photometer. Finally, the LDL levels were determined with formula, $LDL = \text{Total cholesterol} - HDL - 1/5 \text{ Triglyceride}$.

Then these chickens were slaughtered and bled. Then, the visceral organs (liver, spleen, thymus, and bursa), breast muscles, and thigh muscles were collected; and the percentage of offal was calculated.

Antioxidant Status

The right thigh muscles of slaughtered poultry were separated from the carcasses, washed in 0.9% NaCl solution, individually packaged, and immediately frozen at -20°C until analysis. Frozen samples were thawed, and 1g of each sample was diluted with 9mL of ice-cold phosphate-buffered saline (pH=7.2), homogenized in a homogenizer, and centrifuged at 7378g for 15min. Malondialdehyde levels in the meat tissue homogenate supernatant and serum were tested using a commercial test kit (E-BC-K025-S; Elabscece).

Digestibility

Eight cages were randomly selected at 30 days of age, with two birds per cage transferred to individual cages for a period (31-40 days) for digestibility testing following the procedure (Solangi et al. 2020). The adaptation period was given for three days and continued with measurements of feed intake, water intake, and total feces expelled for six days (age 34-39 d). The feces collected in 24 hours were weighed, and samples were taken from each chicken and then dried at 60°C, followed by an analysis of crude protein, ether extract and crude fiber content (AOAC 2005). The same analysis was also carried out on the feed samples.

Statistics

The study data were tabulated and started with the normality test with SPSS (SPSS Inc., Chicago, Illinois) using the Shapiro-Wilk test. Then one-way ANOVA test was carried out to see the effect of the treatment. Orthogonal polynomial contrast analysis was used to analyze the linear and quadratic effect of the level of CGE addition in the treatment diets.

RESULTS

Growth Performance of Broiler

Table 3 shows the effect of adding CGE on broiler growth performance. The growth of broiler body weight in the starter period (age 1-20 d), finisher period (age 21-40 d), and overall period (age 1-40 d) increased linearly ($P=0.022$), ($P=0.007$), and ($P=0.004$) with an increase in the level of addition of CGE from 0, 0.1, 0.2 and 0.3g/kg in the diet. Feed intake was not affected by the addition of CGE in all measured phases. Feed conversion was not significant in the starter phase (age 1-20 d) but decreased linearly in the finisher and overall phases ($P=0.026$ and $P=0.017$) due to the addition of CGE in the diets.

Carcass Characteristics

The effect of the addition of CGE in broiler diets on carcass characteristics is shown in Table 4. The addition

of CGE in the diet linearly increased the percentage of thigh and drumstick ($P=0.03$) and decreased abdominal fat ($P=0.02$) but did not affect the percentage of eviscerated yield, breast, liver, gizzard, pancreas, gallbladder, heart and spleen.

Table 1: Polyphenol content of catechin gambir extract

| Item | Value (%) |
|--|-----------|
| Catechin | 95.87 |
| Total Phenol | 45.98 |
| Condensed Tanin | 1.98 |
| Inhibition percentage (IC ₅₀), (µg/mL) | 93.92 |

Value are means, n=4

Table 2: Analyzed nutrient composition g/kg of commercial diets

| Item | Pre-Starter (1-10 d) | Starter (11-20 d) | Finisher (21-40 d) |
|--|----------------------|-------------------|--------------------|
| Dry matter, g/kg | 884 | 879 | 878 |
| Crude protein, g/kg | 227.21 | 212.40 | 188.12 |
| Lysine, g/kg | 12 | 11 | 10 |
| Methionine, g/kg | 6 | 5 | 4 |
| Ether extract, g/kg | 25 | 38 | 41 |
| Crude fiber, g/kg | 28 | 30 | 43 |
| Ash, g/kg | 58 | 59 | 58 |
| Calcium, g/kg | 10 | 9 | 8 |
| Phosphorus, g/kg | 6 | 5 | 4 |
| Apparent metabolizable energy, Kcal/kg | 2.900 | 3.000 | 3.100 |

Plasma Constituents

The addition of CGE in the diet to blood plasma constituents of broilers (Table 5) shows that CGE linearly decreased total cholesterol ($P=0.036$), decreased low-density lipoprotein cholesterol ($P=0.034$), increased high-density lipoprotein cholesterol ($P=0.017$), and increased total protein ($P=0.001$) in the blood plasma of broilers aged 40 days. However, the content of Triglyceride and glucose in blood plasma was not affected by eating CGE.

Antioxidant Status

The effect of the addition of CGE in the diet on the antioxidant status of broilers aged 40 days is shown in Table 6. Antioxidant activity as measured by the concentration of MDA was found that increasing levels of CGE in broiler diets linearly decreased the concentration of MDA in serum ($P=0.012$) and tissue thigh meat ($P=0.001$) in broilers aged 40 days.

Nutrient Digestibility

Nutrient digestibility in broilers fed diets with the addition of CGE is shown in Table 6. The addition of CGE in diets linearly increased crude protein digestibility ($P=0.037$) and decreased ether extract digestibility ($P=0.006$). However, it did not affect crude fiber digestibility measured in broilers at 31-40 days of age.

DISCUSSION

The rapid growth of broilers with high feed efficiency is the goal of the broiler industry in producing healthy broiler meat. In this study, CGE was used as a source of catechins in broiler feed which acts as a herbal feed additive. In the pre-starter phase (age 1-10 d), no additional CGE was added to the diets to achieve average growth according to breeder standards, namely 320g with an FCR of 1.03 (JCI 2019).

Table 3: Effects of dietary CGE supplement on the growth performance of broiler

| Item | CGE in basal diet, g/kg | | | | SEM | P-value | |
|----------------------------------|-------------------------|-------|-------|-------|-------|---------|-----------|
| | 0 | 0.1 | 0.2 | 0.3 | | Linear | Quadratic |
| Initial body weight, 10 d of age | 239.8 | 238.7 | 241.9 | 243.5 | 2.43 | - | - |
| Starter period, 11-20 d of age | | | | | | | |
| Weight gain, g | 586.7 | 590.8 | 654.9 | 626.6 | 27.18 | 0.022 | 0.718 |
| Feed intake, g | 803.7 | 796.9 | 784.9 | 789.6 | 12.72 | 0.282 | 1.256 |
| Feed conversion | 1.37 | 1.35 | 1.21 | 1.26 | 0.05 | 0.053 | 0.534 |
| Finisher period, 21-40 d of age | | | | | | | |
| Weight gain, g | 1736 | 1692 | 1766 | 1854 | 24.48 | 0.007 | 0.035 |
| Feed intake, g | 3350 | 3322 | 3271 | 3162 | 74.22 | 0.075 | 0.860 |
| Feed conversion | 1.93 | 1.97 | 1.85 | 1.71 | 0.06 | 0.026 | 0.096 |
| Overall, 1-40 d of age | | | | | | | |
| Weight gain, g | 2562 | 2521 | 2663 | 2724 | 28.38 | 0.004 | 0.080 |
| Feed intake, g | 4434 | 4396 | 4330 | 4227 | 84.60 | 0.078 | 1.709 |
| Feed conversion | 1.73 | 1.74 | 1.63 | 1.55 | 0.04 | 0.017 | 0.199 |

SEM=Standard error mean; P=Probability of significant.

Table 4: Effect of catechin gambir extract on carcass characteristics at 40 day

| Response parameter, % of total weight | CGE in basal diet, kg/kg | | | | SEM | P-value | |
|---------------------------------------|--------------------------|-------|-------|-------|------|---------|-----------|
| | 0 | 0.1 | 0.2 | 0.3 | | Linear | Quadratic |
| Eviscerated yield | 70.23 | 69.97 | 69.93 | 69.85 | 0.14 | 0.077 | 0.621 |
| Breast | 22.31 | 24.00 | 23.78 | 23.62 | 0.72 | 5.240 | 0.064 |
| Thigh and drumstick | 17.98 | 18.30 | 18.76 | 21.50 | 0.89 | 0.030 | 0.130 |
| Liver | 2.03 | 2.10 | 1.99 | 1.92 | 0.06 | 0.085 | 0.187 |
| Gizzard | 1.87 | 1.89 | 1.90 | 1.72 | 0.05 | 0.058 | 0.054 |
| Abdominal fat | 1.08 | 0.99 | 0.96 | 0.91 | 0.02 | 0.020 | 0.095 |
| Pancreas | 0.15 | 0.17 | 0.15 | 0.18 | 0.01 | 0.285 | 1.741 |
| Gallbladder | 0.07 | 0.09 | 0.07 | 0.10 | 0.01 | 0.167 | 1.420 |
| Heart | 0.56 | 0.51 | 0.52 | 0.48 | 0.02 | 0.056 | 0.378 |
| Lymph | 0.14 | 0.16 | 0.14 | 0.16 | 0.01 | 0.259 | 1.677 |

SEM=Standard error mean; P=Probability of significant.

Table 5: Effects of dietary CGE on plasma constituents of broiler chickens at 40 day

| Item | CGE in basal diet, g/kg | | | | SEM | P-value | |
|--------------------------|-------------------------|--------|--------|--------|------|---------|-----------|
| | 0 | 0.1 | 0.2 | 0.3 | | Linear | Quadratic |
| Triglyceride, mg/dL | 57.77 | 58.36 | 50.42 | 53.94 | 2.86 | 0.142 | 1.057 |
| Total Cholesterol, mg/dL | 150.22 | 144.13 | 124.42 | 125.17 | 8.01 | 0.036 | 1.411 |
| LDL, mg/dL | 77.53 | 78.49 | 71.89 | 62.38 | 4.28 | 0.034 | 0.171 |
| HDL, mg/dL | 46.80 | 48.55 | 48.88 | 65.16 | 3.23 | 0.017 | 0.050 |
| Glucose, mg/dL | 136.98 | 131.65 | 121.39 | 122.19 | 0.86 | 0.056 | 0.152 |
| Total protein, mg/dL | 2.57 | 3.83 | 3.68 | 3.51 | 0.12 | 0.001 | 0.012 |

SEM=Standard error mean; P=Probability of significant.

Table 6: Effects of dietary CGE on antioxidant status of broiler chickens at 40 day

| Item | CGE in basal diet, g/kg | | | | SEM | P-value | |
|---|-------------------------|-------|-------|-------|------|---------|-----------|
| | 0 | 0.1 | 0.2 | 0.3 | | Linear | Quadratic |
| Malondialdehyde in blood serum nmol/mL | 4.36 | 4.79 | 3.88 | 3.97 | 0.10 | 0.012 | 0.086 |
| Malondialdehyde in thigh meat tissue nmol/g meat tissue | 15.24 | 15.09 | 12.33 | 10.20 | 0.17 | 0.001 | 0.001 |
| nmol/mg protein | 1.65 | 1.63 | 1.33 | 1.10 | 0.02 | 0.001 | 0.001 |

SEM=Standard error mean; P=Probability of significant.

Table 7: Nutrient Digestibility

| Item | CGE in basal diet, g/kg | | | | SEM | P-value | |
|---------------|-------------------------|-------|-------|-------|------|---------|-----------|
| | 0 | 0.1 | 0.2 | 0.3 | | Linear | Quadratic |
| Crude Protein | 706.1 | 764.5 | 773.4 | 777.5 | 1.27 | 0.037 | 0.078 |
| Ether Extract | 733.4 | 717.4 | 695.0 | 692.5 | 1.01 | 0.006 | 0.037 |
| Crude Fiber | 327.5 | 295.0 | 330.0 | 322.5 | 0.82 | 0.862 | 0.113 |

SEM=Standard error mean; P=Probability of significant.

The addition of CGE 0.3g/kg in the basal diet from the starter to the finisher period (age 10-40 d) resulted in a 6.30% increase in body weight gain and a 1.61% improvement in FCR compared to the control. In line with this, a 10% increase in broiler growth and improved feed conversion was also reported with green tea catechins at the level of 0.3g/kg (El-Hack et al. 2020). Catechins are

classified as flavonoid compounds that positively affect broiler performance at specific doses (Viveros et al. 2011). The results of a review of 78 research articles (Prihambodo et al. 2021) show that the addition of 0.8g/kg flavonoid compounds in the diet could increase broiler daily body weight gain (ADG) and improve FCR in the finisher phase but had no effect on the starter phase. In this study, the

concentration of catechin compounds in the diets of 0.287g/kg (treatment 0.3g/kg CGE) got a positive response to body weight gain and improvement in FCR. However, (Chamorro et al. 2013) reported a negative response of catechin compounds on broiler performance, starting to be seen at the catechin level of 0.42g/kg (5g/kg of Grape seed extracts).

Catechins act as antioxidant compounds. The antioxidant activity was seen from the IC₅₀ value is 93.92g/mL (Table 1), indicating that CGE is classified as a potent antioxidant compound (Yeni et al. 2014). Antioxidant compounds have antibacterial activity against various microbial species in the digestive tract, both pathogenic (*Clostridium*) (Abolfathi et al. 2019) and other nuisance bacteria (Srividya et al. 2010). On the other hand, it can increase the population of *Lactobacillus* and *Enterococcus* probiotic bacteria in the intestine (Viveros et al. 2011). Therefore, adding catechins will result in healthier chicken intestines with increased digestibility and absorption capacity (Chamorro et al. 2019). This study also found that crude protein digestibility increased linearly with increasing levels of CGE in the diet (Table 6), so it was reflected in increased growth performance and improved feed conversion (Table 3).

The addition of CGE in the diets increased thigh and drumstick and a tendency to increase breast weight. These results align with some previous reports (Kamboh and Zhu 2013). This increase occurred due to the reduced fat content in broiler meat due to adding phenolic compounds to the diet (Giannenas et al. 2018). The addition of phenolic compounds has an antioxidant effect that can reduce the ratio of n-6/n-3 fatty acids in broiler thigh meat, positively affecting consumer health, especially warding off cardiovascular disease (Saleh et al. 2018).

The increased CGE level in diets linearly decreased the abdominal fat of the broiler. Indeed phenolic compounds generally cause a decrease in abdominal fat (Saracae et al. 2014). Regression analysis of 55 articles on the effect of flavonoids in broiler diets on abdominal fat shows an intercept of 1.65 with a solve of -0.93 (Prihambodo et al. 2021). This decrease in abdominal fat levels is closely related to the effect of catechins in inhibiting fat anabolism and stimulating lipid catabolism (Huang et al. 2015).

Cholesterol is a substance that is produced naturally from the process of fat metabolism in the body, which often binds to protein to form lipoproteins (High-Density Lipoprotein (HDL) and Low-Density Lipoprotein (LDL)). This study found that the addition of CGE in broiler diets reduced total cholesterol and LDL. On the other hand, it increased HDL and total protein (Table 5). The same results were also reported by previous researchers (Saracae et al. 2016) with green tea powder as a source of catechin compounds. These results are closely related to the ability of catechin compounds to reduce enzyme activity to synthesize fatty acids in the liver (Fatty acid synthase and acetyl CoA carboxylase) (Huang et al. 2015).

MDA, a metabolite derived from lipid peroxidation, has been widely used to indicate oxidative damage (Chen et al. 2021). The lower the MDA concentration, the higher the capacity of broiler chickens to clean oxygen free radicals, namely reactive oxygen species (ROS) (Daramola, 2019). Decreased MDA levels in serum and thigh meat tissue may be related to (i) catechin scavenging activity of superoxide and hydroxyl radicals; (ii) chelation

with metal ions and formation of inactive complexes; (iii) stimulation of the synthesis of endogenous antioxidant enzymes in cells (El-Hack et al. 2020).

Adding CGE to the diet can increase the digestibility of crude protein in broilers, which also occurs in the addition of other flavonoid compounds, as reported by Abolfathi et al. (2019); Solangi et al. (2020) and Liu et al. (2020). The content of flavonoids in broiler diets can increase the VH: CD (Villus height: Crypt depth) ratio in the duodenum, jejunum, and ileum (Prihambodo et al. 2021), which indicates a healthier chicken small intestine and higher absorption capacity (Abolfathi et al. 2019). Furthermore, as natural antioxidant compounds, catechins can reduce intestinal oxidative injury in broilers by increasing antioxidant capacity and inhibiting the intestine's inflammatory response, thereby improving intestinal function in digestion and absorption (Song et al. 2019). At a higher dose of catechins at 0.42g/kg (5g/kg grape seed extract), the digestibility of crude protein and essential amino acids (Arginine, Histidine, and Phenylalanine) and non-essential amino acids (Cystine, Glutamic Acid, Glycine, and Proline) began to decrease. Due to the interaction of reactive hydroxyl groups of polyphenols with protein carbonyl groups forming bonds (Chamorro et al. 2013). The decrease in fat digestibility obtained in this study further strengthens that catechins and flavonoids harm fat digestion (Solangi et al. 2020). *In vitro* studies have shown that epigallocatechin gallate interferes with the emulsification, digestion, and dissolution of lipid micelles, which causes a decrease in the absorption of dietary fat in the intestine (Koo and Noh 2007).

13 Conclusion

Based on these results, it can be concluded that CGE can be used as an herbal feed additive for broilers. The addition of 0.3g/kg CGE gave the best results for increasing growth performance, carcass characteristics, blood plasma constituents, antioxidant status, and nutrient digestibility in broilers. Catechins are classified as flavonoids with potent antioxidants activity to provide increased growth performance followed by an increase in carcass characteristics, a decrease in total cholesterol and LDL, an increase in HDL, and antioxidant status as a reflection of an increase in protein digestibility and a decrease in fat digestibility in broilers.

Acknowledgments

The authors are grateful to the Center for Research and Community Service of the Politeknik Pertanian Negeri Payakumbuh, which has facilitated the funding and implementation of the Research Program (PNBP), contract number # 831/PL25/PG/2020.

Author's Contribution

Ramaiyulis supervised the experiment and wrote the original manuscript. Salvia, N. Fati, and T. Malvin conducted the experiment and data analysis. Mairizal prepared tables and finalized draft. The final version of the manuscript was read and approved by all authors.

REFERENCES

Abolfathi ME, Tabeidian SA, Foroozandeh Shahraki AD, Tabatabaei SN and Habibiyan M, 2019. Comparative effects

- of n-hexane and methanol extracts of elecampane (*Inula helenium* L.) rhizome on growth performance, carcass traits, feed digestibility, intestinal antioxidant status and ileal microbiota in broiler chickens. *Archives of Animal Nutrition* 73: 88–110. <https://doi.org/10.1080/1745039X.2019.1581027>
- AOAC, 2005. Official methods of analysis association of official agriculture chemist. Horwitz E (ed). Association of Official Analytical Chemists. 18th Ed. Gaithersburg, MD, USA
- Chamorro S, Romero C, Brenes A, Sánchez-Patán F, Bartolomé B, Viveros A and Arijia I, 2019. Impact of a sustained consumption of grape extract on digestion, gut microbial metabolism and intestinal barrier in broiler chickens. *Food and Function* 10: 1444–1454. <https://doi.org/10.1039/c8fo02465k>
- Chamorro, Viveros A, Centeno C, Romero C, Arijia I and Brenes A, 2013. Effects of dietary grape seed extract on growth performance, amino acid digestibility and plasma lipids and mineral content in broiler chicks. *Animal* 7: 555–561. <https://doi.org/10.1017/S1751731112001851>
- Chen R, Yang M, Song YD, Wang RX, Wen C, Liu Q, Zhou YM and Zhuang S, 2021. Effect of anhydrous betaine and hydrochloride betaine on growth performance, meat quality, postmortem glycolysis, and antioxidant capacity of broilers. *Poultry Science* 101: 687. <https://doi.org/10.1016/j.psj.2021.101687>
- Chun OK, Kim DO and Lee CY, 2003. Superoxide Radical Scavenging Activity of the major polyphenols in fresh plums. *Journal of Agricultural and Food Chemistry* 51: 8067–8072. <https://doi.org/10.1021/jf034740d>
- Daramola OT, 2019. Medicinal plants leaf meal supplementation in broiler chicken diet: Effects on performance characteristics, serum metabolite and antioxidant status. *Animal Research International* 16: 3334–3342.
- El-Hack MEA, Elnesr SS, Alagawany M, Gado A, Noreldin AE and Gabr AA, 2020. Impact of green tea (*Camellia sinensis*) and epigallocatechin gallate on poultry. *World's Poultry Science Journal* 76: 49–63. <https://doi.org/10.1080/00439339.2020.1729672>
- Erinle TJ, Oladokun S, MacIsaac J, Rathgeber B and Adewole D, 2022. Dietary grape pomace – effects on growth performance, intestinal health, blood parameters, and breast muscle myopathies of broiler chickens. *Poultry Science* 101: 101519. <https://doi.org/10.1016/j.psj.2021.101519>
- Ferdinal N, 2014. A Simple purification method of catechin from gambir. *International Journal on Advanced Science, Engineering and Information Technology* 4: 53–55.
- Giannenas I, Bonos E, Skoufos I, Tzora A, Stylianaki I, Lazari D, Tsinas A, Christaki E and Florou-Paneri P, 2018. Effect of herbal feed additives on performance parameters, intestinal microbiota, intestinal morphology and meat lipid oxidation of broiler chickens. In *British Poultry Science* 59: 545-553. Taylor and Francis. <https://doi.org/10.1080/00071668.2018.1483577>
- Gungor E, Altop A and Erener G, 2021. Effect of raw and fermented grape seed on growth performance, antioxidant capacity, and cecal microflora in broiler chickens. *Animal* 15: 100194. <https://doi.org/10.1016/j.animal.2021.100194>
- Huang JB, Zhang Y, Zhou YB, Wan XC and Zhang JS, 2015. Effects of epigallocatechin gallate on lipid metabolism and its underlying molecular mechanism in broiler chickens. *Journal of Animal Physiology and Animal Nutrition* 99: 719–727. <https://doi.org/10.1111/jpn.12276>
- JCI, 2019. Broiler. *Japfa Comfeed Indonesia*, 2 P. www.japfacomfeed.co.id
- Kamboh AA and Zhu WY, 2013. Effect of increasing levels of bioflavonoids in broiler feed on plasma anti-oxidative potential, lipid metabolites, and fatty acid composition of meat. *Poultry Science* 92: 454–461. <https://doi.org/10.3382/ps.2012-02584>
- Kassim MJ, Hussin MH, Achmad A, Dahon NH, Suan TK and Hamdan HS, 2011. Determination of total phenol, condensed tannin and flavonoid contents and antioxidant. *Majalah Farmasi Indonesia* 22: 50–59.
- Koo SI and Noh SK, 2007. Green tea as inhibitor of the intestinal absorption of lipids: potential mechanism for its lipid-lowering effect. *Journal of Nutritional Biochemistry* 18: 179–183. <https://doi.org/10.1016/j.jnutbio.2006.12.005>
- Lillehoj H, Liu Y, Calsamiglia S, Fernandez-Miyakawa ME, Chi F, Cravens RL, Oh S and Gay CG, 2018. Phytochemicals as antibiotic alternatives to promote growth and enhance host health. *Veterinary Research* 49: 1–19. <https://doi.org/10.1186/s13567-018-0562-6>
- Liu HS, Mahfuz SU, Wu D, Shang QH and Piao XS, 2020. Effect of chestnut wood extract on performance, meat quality, antioxidant status, immune function, and cholesterol metabolism in broilers. *Poultry Science* 99: 4488–4495. <https://doi.org/10.1016/j.psj.2020.05.053>
- Prihambodo TR, Sholikin MM, Qomariyah N, Jayanegara A, Batubara I, Utomo DB and Nahrowi N, 2021. Effects of dietary flavonoids on performance, blood constituents, carcass composition and small intestinal morphology of broilers: A meta-analysis. *Animal Bioscience* 34: 434–442. <https://doi.org/10.5713/ajas.20.0379>
- Saleh H, Golian A, Kermanshahi H and Mirakzehi MT, 2018. Antioxidant status and thigh meat quality of broiler chickens fed diet supplemented with α -tocopherolacetate, pomegranate pomace and pomegranate pomace extract. *Italian Journal of Animal Science* 17: 386–395. <https://doi.org/10.1080/1828051X.2017.1362966>
- Saraee MHA, Seidavi A, Dadashbeiki M and Tufarelli V, 2014. Effect of dietary supplementation with different levels of green tea powder and fish oil or their combination on carcass characteristics in broiler chickens. *Pakistan Journal of Zoology* 46: 1767–1773.
- Saraei MH, Seidavi AR, Dadashbeiki M and Edens FW, 2016. Response of plasma constituents and body measurement in broiler chickens fed fish oil and green tea powder. *Archivos de Medicina Veterinaria* 48: 61–68. <https://doi.org/10.4067/S0301-732X2016000100008>
- Solangi MAAMK, Mughal GA, Leghari IH, Rajput MN, Khaskheli AA, Korejo TA, Rajput NA, Memon QA and Kumbhar MA, 2020. Influence of *Camellia sinensis* on nutrients digestibility in broiler. *Pesquisa Agropecuaria Brasileira* 9: 1325–1333. <https://doi.org/10.19045/bspab.2020.90138>
- Song J, Lei X, Luo J, Everaert N, Zhao G, Wen J and Yang Y, 2019. The effect of Epigallocatechin-3-gallate on small intestinal morphology, antioxidant capacity and anti-inflammatory effect in heat-stressed broilers. *Journal of Animal Physiology and Animal Nutrition* 103: 1030–1038. <https://doi.org/10.1111/jpn.13062>
- Srividya A, Dhanabal S, Misra V and Suja G, 2010. Antioxidant and antimicrobial activity of *Alpinia officinarum*. *Indian Journal of Pharmaceutical Sciences* 72: 145–148.
- Viveros A, Chamorro S, Pizarro M, Arijia I, Centeno C and Brenes A, 2011. Effects of dietary polyphenol-rich grape products on intestinal microflora and gut morphology in broiler chicks. *Poultry Science* 90: 566–578. <https://doi.org/10.3382/ps.2010-00889>
- Widiyarti G, Sundowo A, Filaila E and Laksmono AJ, 2020. The Mechanically Extraction Process of Gambir (*Uncaria gambir Roxb*) from Limapuluh Kota, West Sumatera and Its Antioxidant activity. *The Journal of Pure and Applied Chemistry Research* 9: 8–15. <https://doi.org/10.21776/ub.jpacr.2020.009.01.509>
- Yeni G, Syamsu K, Suparno O, Mardiyati E and Muchtar H, 2014. Repeated extraction process of raw gambirs (*Uncaria gambir Roxb*) for the catechin production as an antioxidant. *International Journal of Applied Engineering Research* 9: 24565–24578.

Scopus

ORIGINALITY REPORT

21%
SIMILARITY INDEX

19%
INTERNET SOURCES

19%
PUBLICATIONS

2%
STUDENT PAPERS

PRIMARY SOURCES

| | | |
|----------|--|-----------|
| 1 | www.tandfonline.com Internet Source | 3% |
| 2 | indonesianjpharm.farmasi.ugm.ac.id Internet Source | 2% |
| 3 | www.animbiosci.org Internet Source | 1% |
| 4 | www.ijvets.com Internet Source | 1% |
| 5 | www.researchgate.net Internet Source | 1% |
| 6 | mafiadoc.com Internet Source | 1% |
| 7 | "Fundamental and Applied Scientific Research in the Development of Agriculture in the Far East (AFE-2021)", Springer Science and Business Media LLC, 2022 Publication | 1% |
| 8 | Seyyed Javad Hosseini-Vashan, Moreza Safdari-Rostamabad, Ali Hossein Piray, Hadi | 1% |

Sarir. "The growth performance, plasma biochemistry indices, immune system, antioxidant status, and intestinal morphology of heat-stressed broiler chickens fed grape (*Vitis vinifera*) pomace", *Animal Feed Science and Technology*, 2019

Publication

9

www.bdbotsociety.org

Internet Source

1 %

10

E A Zebua, J Silalahi, E Julianti. " Hypoglycemic activity of gambier (.) drinks in alloxan-induced mice ", *IOP Conference Series: Earth and Environmental Science*, 2018

Publication

<1 %

11

www.science.gov

Internet Source

<1 %

12

t-stor.teagasc.ie

Internet Source

<1 %

13

repository.unja.ac.id

Internet Source

<1 %

14

Shun Chen, Huijuan Liu, Jiaqi Zhang, Binbin Zhou, Su Zhuang, Xiaofang He, Tian Wang, Chao Wang. "Effects of different levels of rutin on growth performance, immunity, intestinal barrier and antioxidant capacity of broilers", *Italian Journal of Animal Science*, 2022

Publication

<1 %

| | | |
|----|---|------|
| 15 | www.poultryscience.org Internet Source | <1 % |
| 16 | www.researcherslinks.com Internet Source | <1 % |
| 17 | M. H. Farahat, F. M. Abdallah, H. A. Ali, A. Hernandez-Santana. "Effect of dietary supplementation of grape seed extract on the growth performance, lipid profile, antioxidant status and immune response of broiler chickens", animal, 2016 Publication | <1 % |
| 18 | www.thieme-connect.com Internet Source | <1 % |
| 19 | comum.rcaap.pt Internet Source | <1 % |
| 20 | dalspace.library.dal.ca Internet Source | <1 % |
| 21 | Submitted to Higher Education Commission Pakistan Student Paper | <1 % |
| 22 | www.elis.sk Internet Source | <1 % |
| 23 | H.S. Liu, S.U. Mahfuz, D. Wu, Q.H. Shang, X.S. Piao. "Effect of chestnut wood extract on performance, meat quality, antioxidant status, | <1 % |

immune function, and cholesterol metabolism in broilers", Poultry Science, 2020

Publication

24

www.indianjournals.com

Internet Source

<1 %

25

Alexandra Desbruslais, Alexandra L. Wealleans. "Oxidation in Poultry Feed: Impact on the Bird and the Efficacy of Dietary Antioxidant Mitigation Strategies", Poultry, 2022

Publication

<1 %

26

R. Chen, M. Yang, Y.D. Song, R.X. Wang, C. Wen, Q. Liu, Y.M. Zhou, S. Zhuang. "Effect of anhydrous betaine and hydrochloride betaine on growth performance, meat quality, postmortem glycolysis, and antioxidant capacity of broilers", Poultry Science, 2021

Publication

<1 %

27

jasbsci.biomedcentral.com

Internet Source

<1 %

28

www.medicaljournalofcairouniversity.net

Internet Source

<1 %

29

"Abstract Section", Maturitas, 2003

Publication

<1 %

30

M. Nardoia, C. Romero, A. Brenes, I. Arija, A. Viveros, C. Ruiz-Capillas, S. Chamorro. "Addition of fermented and unfermented

<1 %

grape skin in broilers' diets: effect on digestion, growth performance, intestinal microbiota and oxidative stability of meat", animal, 2019

Publication

31

janimscitechnol.biomedcentral.com

Internet Source

<1 %

32

Norizaty Irman, Nur Hanis Abd Latif, Nicolas Brosse, Francois Gambier, Firda Aulya Syamani, M. Hazwan Hussin. "Preparation and characterization of formaldehyde-free wood adhesive from mangrove bark tannin", International Journal of Adhesion and Adhesives, 2022

Publication

<1 %

33

archive.org

Internet Source

<1 %

34

repository.pertanian.go.id

Internet Source

<1 %

35

www.mdpi.com

Internet Source

<1 %

36

Young Seok Oh. "Singlet Oxygen Quenching Activities of Various Fruit and Vegetable Juices and Protective Effects of Apple and Pear Juices against Hematolysis and Protein Oxidation Induced by Methylene Blue

<1 %

37

core-cms.prod.aop.cambridge.org

Internet Source

<1 %

38

s3-eu-west-1.amazonaws.com

Internet Source

<1 %

39

J. C. Han, X. D. Yang, L. M. Zhang, W. L. Li, T. Zhang, Z. Y. Zhang, J. H. Yao. "Effects of 1慣-Hydroxycholecalciferol and Phytase on Growth Performance, Tibia Parameter and Meat Quality of 1- to 21-d-old Broilers", Asian-Australasian Journal of Animal Sciences, 2009

Publication

<1 %

40

Majid Gholami, Mohammad Chamani, Alireza Seidavi, Ali Asghar Sadeghi, Mehdi Aminafschar. "Effects of stocking density and environmental conditions on performance, immunity, carcass characteristics, blood constituents, and economical parameters of cobb 500 strain broiler chickens", Italian Journal of Animal Science, 2020

Publication

<1 %

41

Mohamed M. Alghirani, Eric Lim Teik Chung, Nafeesa Abu Kassim, Yee Lyn Ong et al. "Effect of Brachiaria decumbens as a novel supplementation on the production

<1 %

performance of broiler chickens", Tropical Animal Health and Production, 2022

Publication

42

S. H. Abu Hafsa, S. A. Ibrahim. "Effect of dietary polyphenol-rich grape seed on growth performance, antioxidant capacity and ileal microflora in broiler chicks", Journal of Animal Physiology and Animal Nutrition, 2018

Publication

<1 %

43

S. Yarmohammadi Barbarestani, V. Jazi, H. Mohebodini, A. Ashayerizadeh, A. Shabani, M. Toghyani. "Effects of dietary lavender essential oil on growth performance, intestinal function, and antioxidant status of broiler chickens", Livestock Science, 2020

Publication

<1 %

44

ikee.lib.auth.gr

Internet Source

<1 %

45

organicplusnet.files.wordpress.com

Internet Source

<1 %

46

rp2u.unsyiah.ac.id

Internet Source

<1 %

47

www.jenschristianholm.dk

Internet Source

<1 %

48

Abdullahi Abdulmajid, Tuan Sherwyn Hamidon, Afidah Abdul Rahim, M. Hazwan Hussin. "Physicochemical studies of tamarind

<1 %

shell tannins as a potential green rust converter", BioResources, 2019

Publication

49

Ahmed M. Fikry, Adel I. Attia, Ismail E. Ismail, Mahmoud Alagawany, Fayiz M. Reda. "Dietary citric acid enhances growth performance, nutrient digestibility, intestinal microbiota, antioxidant status, and immunity of Japanese quails", Poultry Science, 2021

Publication

50

Gengsheng Xiao, Liwei Zheng, Xia Yan, Li Gong, Yang Yang, Qien Qi, Xiangbin Zhang, Huihua Zhang. "Effects of Dietary Essential Oils Supplementation on Egg Quality, Biochemical Parameters, and Gut Microbiota of Late-Laying Hens", Animals, 2022

Publication

51

Jiao Song, Xin Lei, Jingxian Luo, Nadia Everaert, Guiping Zhao, Jie Wen, Ye Yang. "The effect of Epigallocatechin-3-gallate on small intestinal morphology, antioxidant capacity and anti-inflammatory effect in heat-stressed broilers", Journal of Animal Physiology and Animal Nutrition, 2019

Publication

52

Muhammad Abdul Basit, Abdul Kadir Arifah, Teck Chwen Loh, Abdul Aziz Saleha, Annas Salleh, Ubedullah Kaka, Sherifat Banke Idris.

<1 %

<1 %

<1 %

<1 %

"Effects of graded dose dietary supplementation of Piper betle leaf meal and Persicaria odorata leaf meal on growth performance, apparent ileal digestibility, and gut morphology in broilers", Saudi Journal of Biological Sciences, 2020

Publication

53

N. Rajput, S. Ali, M. Naeem, M.A. Khan, T. Wang. "The effect of dietary supplementation with the natural carotenoids curcumin and lutein on pigmentation, oxidative stability and quality of meat from broiler chickens affected by a coccidiosis challenge", British Poultry Science, 2014

Publication

<1 %

54

N. Sahin, M. Onderci, K. Sahin, G. Cikim, O. Kucuk. "Magnesium Proteinate Is More Protective than Magnesium Oxide in Heat-Stressed Quail", The Journal of Nutrition, 2005

Publication

<1 %

55

Nikola Puvača, Vincenzo Tufarelli, Ilias Giannenas. "Essential Oils in Broiler Chicken Production, Immunity and Meat Quality: Review of Thymus vulgaris, Origanum vulgare, and Rosmarinus officinalis", Agriculture, 2022

Publication

<1 %

56

Saeed Safavipour, Sayed Ali Tabeidian, Majid Toghyani, Amir Davar Foroozandeh Shahraki,

<1 %

Gholamreza Ghalamkari, Mahmood Habibian.
"Laying performance, egg quality, fertility,
nutrient digestibility, digestive enzymes
activity, gut microbiota, intestinal
morphology, antioxidant capacity, mucosal
immunity, and cytokine levels in meat-type
Japanese quail breeders fed different
phytogenic levels", Research in Veterinary
Science, 2022

Publication

57

digital.library.adelaide.edu.au

Internet Source

<1 %

58

docplayer.org

Internet Source

<1 %

59

hdl.handle.net

Internet Source

<1 %

60

highcholesterolcure.info

Internet Source

<1 %

61

ir.amu.ac.in

Internet Source

<1 %

62

onlinelibrary.wiley.com

Internet Source

<1 %

63

sciendo.com

Internet Source

<1 %

64

www.european-poultry-science.com

Internet Source

<1 %

65

www.frontiersin.org

Internet Source

<1 %

66

M. Sholikin, S. Sadarman, A. Irawan, T. Prihambodo, N. Qomariyah, A. Wahyudi, J. Nomura, N. Nahrowi, A. Jayanegara. "Antimicrobial peptides as an additive in broiler chicken nutrition: a meta-analysis of bird performance, nutrient digestibility and serum metabolites", *Journal of Animal and Feed Sciences*, 2021

Publication

<1 %

67

Tuty Anggraini, Neswati, Alfi Asben. "Gambir Quality from West Sumatra Indonesia Processed With Traditional Extraction", *IOP Conference Series: Earth and Environmental Science*, 2019

Publication

<1 %

68

"Mediterranean Fruits Bio-wastes", Springer Science and Business Media LLC, 2022

Publication

<1 %

69

"Obesity and Cancer", Springer Science and Business Media LLC, 2021

Publication

<1 %

70

Alberto Barbabosa Pliego, Masoomeh Tavakoli, Ameer Khusro, Alireza Seidavi et al. "Beneficial and adverse effects of medicinal plants as feed supplements in poultry

<1 %

nutrition: a review", Animal Biotechnology,
2020

Publication

71

RP Turcu, M Olteanu, RD Criste, TD Panaite, M Ropotă, PA Vlaicu, D Drăgotoiu. "Grapeseed Meal Used as Natural Antioxidant in High Fatty Acid Diets for Hubbard Broilers", Brazilian Journal of Poultry Science, 2019

Publication

<1 %

72

U. Gadde, W. H. Kim, S. T. Oh, Hyun S. Lillehoj. "Alternatives to antibiotics for maximizing growth performance and feed efficiency in poultry: a review", Animal Health Research Reviews, 2017

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On