




# Plagiarism Checker X - Report

Originality Assessment

# 19%



**Overall Similarity**

**Date:** Mar 16, 2023

**Matches:** 717 / 3794 words

**Sources:** 26

**Remarks:** Low similarity detected, check with your supervisor if changes are required.

**Verify Report:**

[V](#) [r](#) [t](#) [f](#) [c](#) [t](#) [n](#) [l](#) [n](#)

358 To cite this paper: Ramaiyulis R, Metri Y, Irda I, Kurnia D, and Syukriani D (2022). Effects of Tannin-containing Supplement on Enteric Methane Emissions, Total Digestible Nutrient, and Average Daily Gain of Local Indonesian Beef Cattle. World Vet. J., 12 (4): 358-362. DOI: <https://dx.doi.org/10.54203/scil.2022.wvj45> 2022, Scienceline Publication World's Veterinary Journal World Vet J, 12(4): 358-362, December 25, 2022 ISSN 2322-4568

1 Effects of Tannin-containing Supplement on Enteric Methane Emissions, Total Digestible Nutrient, and Average Daily Gain of Local Indonesian Beef Cattle Ramaiyulis

Ramaiyulis1\* , Yurma Metri2 , Irzal Irda1 , Dihan Kurnia1 , and Debby Syukriani1 1Animal Production, Agricultural Polytechnic of Payakumbuh, Harau, Lima Puluh Kota, 26271, Indonesia 2Agriculture Department, Sekolah Tinggi Haji Agussalim, Bukittinggi 26117 Indonesia \*Corresponding author's Email:

ramaiyulis@gmail.com ABSTRACT Reducing methane (CH<sub>4</sub>) emissions is one of the most critical goals in ruminant nutrition.

9 This study aimed to evaluate the effect of concentrate and tannin supplementation on the mitigation of methane gas in Indonesian local beef cattle. The current study was conducted in vivo using 12 Bali cattle using a completely randomized design with four treatments and three replicates. Cattle were 2 fed a basal ration with field grass (control), the addition of concentrate 25% dry matter

(DM) ration no tannin as well as tannin supplemented in concentrate at levels of 0.12% and 0.18% of DM concentrate. The concentrate contains 7.5% crude protein and 71.25% total digestible nutrients and tannin supplementation using gambir (*Uncaria gambir* Indonesia) tannin extract. The parameters measured were apparent digestibility, total digestible nutrients, methane production, 1 and average daily gain. The results showed that

concentrate addition significantly increased DM consumption, crude protein digestibility, and total digestible nutrients. Supplementation of 0.18% tannin in concentrate can mitigate 49.7% methane gas production resulting in energy efficiency, which was reflected in the weight gain rate of 0.75 kg/day. In conclusion, present results suggest that the supplementation of 0.18% gambier tannin extract in concentrate could be considered a suitable feed additive to mitigate methane gas production and increase the average daily

gain of Indonesian local beef cattle. Keywords: Digestibility, Feed supplement, Gambir, Methane, Tannin ORIGINAL ARTICLE pii: S232245682200045-12 Received: 18 October

2022 Accepted: 03 December 2022 INTRODUCTION The population of beef cattle in

Indonesia in 2021 was 18 million heads, increasing about 2.3% since 2016 (BPS, 2021).

The increase in greenhouse gases in the atmosphere from components of carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), and nitrous oxide (N<sub>2</sub>O) is a challenge behind the increase in

livestock population. Methane production in ruminants contributes 56% of total agricultural greenhouse gas emissions and 93% of total livestock emissions globally (Watts et al.,

2021). In addition to impacting the environment as a greenhouse gas, methane emissions are a waste of energy that livestock should use for production. About 3-12% of the gross

energy of feed consumed by ruminants is lost as methane (Ku-Vera et al., 2020).

Strategies to reduce methane emissions are grouped into animal breeding, dietary

manipulation, and rumen manipulation (Arndt et al., 2021). Meta-analysis showed <sup>4</sup> that

the addition of concentrate and tannin supplementation decreased CH<sub>4</sub> Yield (g/kg DMI)

by 13.7-17.4% and CH<sub>4</sub>/Gain by 20.1% (Congio et al., 2021). Thus, in the current study,

the concentrate was added to tannin supplementation from gambir extract (*Uncaria gambir*

Indonesia) to mitigate CH<sub>4</sub> emission. This study hypothesized that adding concentrate

supplemented with tannins could mitigate CH<sub>4</sub> emissions and increase the total digestible

nutrients, thus reflecting an increase in average daily gain. <sup>4</sup> Therefore, this study aimed

to determine the effect of tannin concentrate and supplementation on enteric CH<sub>4</sub>

production, total digestible nutrient, and average daily gain in local Indonesian

cattle. MATERIALS AND METHODS Ethical approval <sup>2</sup> This study was carried out

according to standard protocols without causing discomfort or injury to the cattle.

Furthermore, the experimental procedure was approved by the <sup>18</sup> Center for Research

and Community Service at Agricultural Polytechnic of Payakumbuh, referring to

Government Regulation of the Republic of Indonesia No. 95 of 2012 <sup>23</sup> concerning

veterinary public health and animal welfare. DOI:

<https://dx.doi.org/10.54203/scil.2022.wvj45>

359 To cite this paper: Ramaiyulis R, Metri Y, Irda I, Kurnia D, and Syukriani D (2022). 1

Effects of Tannin-containing Supplement on Enteric Methane Emissions, Total Digestible Nutrient, and Average Daily Gain of Local Indonesian Beef Cattle. World Vet. J., 12 (4):

358-362. DOI: <https://dx.doi.org/10.54203/scil.2022.wvj45> Animal and treatment diets 4

The experiment was conducted in June-September 2021 at the Agricultural Polytechnic of

Payakumbuh, Sumatera Barat, Indonesia. The climatic data referred to in AccuWeather

(2021) recorded throughout the study is the end of summer and 20 the beginning of the

entry of the rainy season with a daily temperature range of 24-34 °C. The study used 12

male Bali cattle (*Bos sondaicus*) of similar aged 1.5-1.8 years, with initial body weights of

237.31±4.85 kg. The cattle were placed in a metabolic cage measuring 1.8×1.5 m. Newly

entered cattle were given the parasite drug Wormectin 1ml/ 50 kg of body weight (Medion,

2022). The animals and treatments were allocated randomly according to 10 a completely

randomized design with four treatments and three replications. The treatments were 100%

basal ration (control), the addition concentrate of 25% dry matter (DM) ration without

tannins (T0), and the addition of 0.12 (T2) and 0.18 (T3) % of DM concentrate gambir

tannin extract. Tannin extract was obtained from the extraction of tannins from the gambir

plant with water as a solvent in the gambier industry according to the method by Fauza

(2014). Basal ration using mixed field grass from the genus *Paspalum* and the concentrate

addition was composed of 30% bran, 20% coconut cake, 30% sago pith, and 20% cassava

with a protein content of 7.5% and a 4 total digestible nutrient (TDN) of 71.25%.

Supplements containing tannins are made 2 with a mixture of 15% brown sugar, 28%

bran, 9% coconut cake, 15% tapioca, 20% soybean meal, 5% urea, 3% salt, 2.5% ultra-

mineral, and 2.5% gambir tannin extract with a crude protein content of 23.69%, and TDN

(78.55%). 21 The chemical composition of the experimental feed is shown in Table 1.

Table 1. Nutritional content of treatment diets Chemical composition Basal diet\* (control)

Concentrate\*\* Level tannin\*\*\* T0 = 0.00 T1 = 0.12 T2 = 0.18 Organic matter (%) 90.33

91.27 90.52 90.14 Crude protein (%) 5.14 6.04 7.65 8.46 Neutral detergent fiber (%) 58.66

52.03 51.09 50.62 Acid detergent fiber (%) 37.85 33.13 32.27 31.84 \*Basal diet: mixed field grass of the Paspalum genus; \*\*add concentrate 25% of dry matter ration; \*\*\*Level tannin (percentage of dry matter concentrate) Sample collection and analysis The experimental period was 40 days, 26 days for the digestibility trial, and 14 days for the daily gain measurement. The digestibility experiment was started with a 10 days preliminary period; the cows were given 100% basal feed to eliminate the effect of the last feed. The next 10 days were the adaptation period to the treatment diet and 6 days to collect digestibility data. The number of rations offered and rejected was weighed to determine the total daily dry matter intake. The excreted feces are collected in a bag tied to the cattle during the collecting period. Then the samples of feed and feces (100 g as fed) were stored in the freezer (-20°C) to be analyzed for their nutritional content. The body weight of cattle was measured at the end of the digestibility data collection as the initial weight and 14 days later as the final weight. The weighing was carried out in the late morning, 16 hours after the previous day's ration was given. After completing the research, the feed and feces samples were dried in an oven at 60°C, then mashed using a 0.5 mm sieve. The concentration of dry matter, organic matter, and crude protein were determined by the method of AOAC (2005), while the fiber fraction was analyzed by the method of Van Soest et al. (1991). Methane gas production was predicted by performing the following Formula 1 according to Jentsch et al. (2007) and Ningrat et al. (2018). Methane (MJ/day = 1.62(DCP) – 0.38(DF) + 3.78(DCF) + 1.49(DN-fe) + 1142 (Formula 1) Where, DCP is a digestible crude protein (g), DF denotes digestible fat (g), DCF signifies digestible crude fiber (g), and DN-fe refers to digestible N-free extract (g) Data analysis The resulting data were tabulated and statistically processed using the statistical package for the social sciences (SPSS, Chicago, USA) was used for analyzing the data, and the one-way ANOVA test was chosen in this software. Duncan's test was chosen to determine the mean significant differences between treatments. The p < 0.05 was considered a significant difference between the groups.

10 RESULTS AND DISCUSSION The effects of supplementation of 25% concentrate (DM diet) and different levels of tannin

supplementation on consumption, digestibility, methane production, and weight gain in Bali cattle are shown in Table 2. As can be seen in Table 2, the addition of concentrate can increase ( $p < 0.05$ ) dry matter consumption, while tannin supplementation does not affect consumption ( $p > 0.05$ ). The addition of concentrate caused a 6.36% reduction in neutral detergent fiber (NDF) in the ration (Control vs. T0-T3, Table 1). The decrease in NDF allows the ration to be digested more quickly in the

360 To cite this paper: Ramaiyulis R, Metri Y, Irda I, Kurnia D, and Syukriani D (2022).

Effects of Tannin-containing Supplement on Enteric Methane Emissions, Total Digestible Nutrient, and Average Daily Gain of Local Indonesian Beef Cattle. World Vet. J., 12 (4):

358-362. DOI: <https://dx.doi.org/10.54203/scil.2022.wvj45>

rumen and accelerates the rate of rumen emptying. The NDF content can reduce dry matter intake (DMI). Souza et al.

(2017) reported a negative correlation between forage NDF and DMI. Condensed tannins

from various plant sources at moderate levels (1-4%) did not have a significant effect on

feed intake (Bunglavan and Duta, 2013). Regression analysis showed that organic

matter digestibility and NDF digestibility had a close relationship with DM consumption ( $p <$

0.05). This relationship showed that dry matter consumption is influenced by the

digestibility of organic matter and NDF. Organic matter and NDF are the main components

of rations in local beef cattle with 100% forage on traditional farms in Indonesia. Organic

matter is all organic components of dietary protein, carbohydrates, fats, and vitamins, and

NDF is a component of cell walls consisting of cellulose, hemicellulose, and lignin

(Lardy, 2018). Table 2. Dry matter intake, apparent digestibility, methane production, and

weight gain measured in Bali beef cattle aged 1.5 years after supplementation of diet with

different levels of tannin

Items	Basal diet	Concentrate	SEM	p value	Level tannin	**
Dry matter intake	2.17b	2.66a	2.77a	2.70a	0.04	<0.01
Metabolic body weight (g/kg)	74.20b	88.35a	93.49a	90.91a	0.09	<0.01
Apparent digestibility (%)	DM 73.74	71.47	71.59	73.02	0.51	0.06
	Organic matter	77.13	75.12			
	75.01	76.35	0.45	0.05		
Crude protein	35.97b	56.30a	55.53a	57.31a	3.00	0.01
Crude fiber						

77.36 76.28 68.89 78.29 2.03 0.05 NDF 66.97 66.59 67.08 67.09 2.83 0.36 ADF 67.55  
60.94 60.17 64.38 1.84 0.05 Cellulose 73.13 69.37 67.20 68.09 1.82 0.08 Total digestible  
nutrients BW (g/kg) 15.45b 19.18a 19.19a 19.06a 0.56 0.01 Percentage of dry  
matter intake 67.89b 72.77a 72.02a 75.35a 0.01 0.04 Methane production (MJ/day)\*  
10.06a 10.40a 9.79a 8.41b 0.15 <0.01 ADG (kg/d) 0.37c 0.47bc 0.59b 0.75a 0.04 0.013

abcmean data with different lowercase letters on the same row show significantly different  
( $p < 0.05$ ). SEM: standard error mean, BW: body weight, DM: Dry matter, BW: Body

weight, NDF: Neutral detergent fiber, ADF: Acid detergent fiber, MJ: Megajoule, TDN:  
Total digestible nutrient ADG: Average daily gain. \*Methane is predicted using the method

of Jentsch et al. (2007); \*\*Percentage of DM concentrate The addition of concentrate and  
supplementation of tannins had no effect ( $p > 0.05$ ) on apparent digestibility except for  
crude protein. The digestibility of crude protein increased with tannin supplementations ( $p$   
< 0.05). In the current study, protein digestibility increased due to an increase in the  
protein level of the diet after the addition of concentrates and tannin supplementation.

Basal diets using tropical forages had a crude protein content of 5.14%, a value  
considered limiting adequate rumen microbial activity, implying sub-optimal conditions in  
the rumen (Sampaio et al., 2010). Rira et al. (2022) reported that condensed tannins did  
not affect feed degradability but interfered with microbial colonization; tannin-rich feeds had  
a relatively lower abundance of fibrinolytic microbes, especially *Fibrobacter* spp.

Furthermore, it is supported by the report of Souza et al. (2010) that the protein content  
below 7 g/kg DM in the basal ration limits the degradation of rumen microorganisms. The  
same finding was also reported by van Kuijk et al. (2022). Tannin compounds can bind  
strongly to proteins (Le Bourvellec and Renard, 2018) which causes a decrease in crude  
protein digestibility in the rumen. According to Ramaiyulis (2021), diet containing 0.12%  
tannin can reduce protein degradation in the rumen from 1.17 to 0.99% per hour in vitro.  
However, in the present study, the opposite results were obtained due to the in vivo nature  
of the study. The tannin-protein complexes formed in the rumen (pH 3.57) make the protein  
resistant to rumen microbial degradation. However, these complexes should dissociate in

the pH environment of the duodenum and proximal abomasum (Getachew et al., 2006). Some studies also indicated the effect of tannins in reducing crude protein degradation in the rumen. However, it did not affect total post-rumen digestibility (Jolazadeh et al., 2015; Canadianti et al., 2020). Total digestible nutrients showed an increase ( $p < 0.05$ ) due to the addition of concentrate and were not affected by tannin supplementation ( $p > 0.05$ ). The addition of concentrate plays a role in supplying additional nutrients because the basal feed is classified as low-quality grass (basal feed with crude protein  $< 8\%$  and NDF  $> 50\%$ ). Rufino et al. (2016) stated that low-quality basal feed requires crude protein supplementation <sup>4</sup> to meet the needs of rumen microbes to develop. The concentrates contain essential nutrients for rumen microbial growth, such as soluble carbohydrates, crude protein sources from urea (non-protein nitrogen), and macro and micro minerals. The fulfillment of rumen microbial nutrition can increase microbial activity to digest feed in the rumen (Ramaiyulis et al., 2019). Table 2 shows that methane production per head of livestock ranged from 8.41 to 10.40 (MJ/d). Methane production is suppressed with 0.18% tannin supplementation ( $p < 0.05$ ). Previous studies also found a 44-49% decrease in the proportion of methane <sup>1</sup> with the addition of supplements containing tannins in forage or fermented straw basal feed

361 To cite this paper: Ramaiyulis R, Metri Y, Irdi I, Kurnia D, and Syukriani D (2022).

Effects of Tannin-containing Supplement on Enteric Methane Emissions, Total Digestible Nutrient, and Average Daily Gain of Local Indonesian Beef Cattle. World Vet. J., 12 (4):

358-362. DOI: <https://dx.doi.org/10.54203/scil.2022.wvj45> (Liu et al., 2019; Ramaiyulis et al., 2021). A dairy cattle diet containing condensed and hydrolyzed tannin compounds can be considered a sustainable approach to reducing the environmental impact of rumination because the feed containing tannins can reduce enteric CH<sub>4</sub> emissions (Stewart et al., 2019). The ratio of Methane/TDN decreased simultaneously <sup>1</sup> with the addition of supplements 0.12 and 0.18 % tannins ( $p < 0.05$ ). Mitigation of methane gas is energy efficient, considering that 5-15% of energy is lost <sup>26</sup> in the form of methane gas during the



rumen fermentation process (Wanapat and Kang, 2015). This efficiency was reflected in the increase of body weight gain in beef cattle from 0.37kg/d (control) to 0.75kg/d <sup>1</sup> with the addition of concentrate and 0.18% tannin supplementation. CONCLUSION The supplemented concentrate with gambir extract tannins can mitigate CH<sub>4</sub> emissions and increase total digestible nutrients. The highest CH<sub>4</sub> mitigation (49.7%) occurred in the basal diets of field grass <sup>1</sup> with the addition of concentrate (75:25% DM) and 0.18% tannin supplementation. The highest CH<sub>4</sub> mitigation can lead to the optimal weight gain of 0.75 kg/d in local Indonesian cattle. DECLARATIONS Acknowledgments The authors are grateful to the <sup>18</sup> Center for Research and Community Service of the Agricultural Polytechnic of Payakumbuh, which has facilitated the funding and implementation of the Research Program (PNBP), contract Number: 2188/PL25/PG/2021. Authors' contribution Ramaiyulis supervised the experiment and wrote the original manuscript. Devi Kumala Sari, Dihan Kurnia, Debby Syukriani, and Irzal Irda conducted the experiment and data analysis. Yurma Metri prepared tables and finalized draft. All authors have read and agreed to the data analysis and the final version of the manuscript. Competing interests <sup>24</sup> The authors declare that they have no conflicts of interest concerning the work presented in this report. Ethical considerations The authors carefully examined all ethical issues concerning plagiarism, approval to publish, errors in fabrication, double publication, and submission. REFERENCES AccuWeather (2021). Weather Padang, West Sumatra Indonesia. Available at: <https://www.accuweather.com/id/id/padang/206120/june-weather/206120?year=2021> Arndt C, Hristov AN, Price WJ, Maclelland SC, Pelaez AM, Cueva SF, Oh J, Bannink A, Bayat AR, Crompton LA et al. (2021). Strategies to mitigate enteric methane emissions by ruminants – A way to approach the 2.0°C target. CABI., Wallingford, UK, pp. 1-53. DOI: <https://www.doi.org/10.31220/agriRxiv.2021.00040> Association of official analytical chemists (AOAC) (2005). <sup>16</sup> Official methods of analysis association of official agriculture chemis. p. 125 In: W. Horwitz (Editor), Association of official analytical chemists, 18th Edition. Gaithersburg, MD, USA, pp. 125-141. Available at: <https://www.worldcat.org/title/official-methods-of-analysis-of-aoac->

international/oclc/62751475 Badan pusat statistik (BPS) (2021). Peternakan dalam Angka 2021. Jakarta, Indonesia. Available at:

[https://www.bps.go.id/publication/2022/07/01/f47af5c5d24ff60405106953/peternakan-](https://www.bps.go.id/publication/2022/07/01/f47af5c5d24ff60405106953/peternakan-dalam-angka-2021.html)

[dalam-angka-2021.html](https://www.bps.go.id/publication/2022/07/01/f47af5c5d24ff60405106953/peternakan-dalam-angka-2021.html) Bunglavan SJ and Dutta N (2013). The use of tannins as organic protectants of proteins in digestion of ruminants. *Journal of Livestock Science*, 4: 67-77.

Available at: [http://livestockscience.in/wp-content/uploads/Buglavan\\_tannin.pdf](http://livestockscience.in/wp-content/uploads/Buglavan_tannin.pdf) Canadianti

M, Yusiati LM, Hanim C, Widyobroto BP, and Astuti A (2020). **11** [The effect of nutmeg leaves tannin \(\*Myristica fragrans\* Houtt\) as protein protecting agents on in vitro nutrient digestibility](#). *Buletin Peternakan*, 44(1): 10-14. DOI:

<https://www.doi.org/10.21059/buletinpeternak.v44i1.47976> Congio GF de S, Bannink A, Mogollón OL, Jaurena G, Gonda H, Gere JI, Cerón-Cucchi ME, Ortiz-Chura A, Tieri M P,

Hernández O et al. (2021). **2** [Enteric methane mitigation strategies for ruminant livestock systems in the Latin America and Caribbean region: A meta-analysis](#). *Journal of Cleaner Production*, 312: 127693. DOI: <https://www.doi.org/10.1016/j.jclepro.2021.127693> Fauza H

(2014). Gambier : Indonesia leading commodities in the past. *International Journal on Advanced Science, Engineering and Information Technology*, 4(6): 67-72. DOI:

<https://www.doi.org/10.18517/ijaseit.4.6.463> Getachew G, Depeters EJ, Pittroff W, Putnam DH, and Dandekar AM (2006). Review: **17** [Does protein in alfalfa need protection from rumen microbes? The Professional Animal Scientist](#), 22(5): 364-373. DOI:

[https://www.doi.org/10.15232/S1080-7446\(15\)311293](https://www.doi.org/10.15232/S1080-7446(15)311293)

362 To cite this paper: Ramaiyulis R, Metri Y, Irda I, Kurnia D, and Syukriani D (2022). **1** [Effects of Tannin-containing Supplement on Enteric Methane Emissions, Total Digestible Nutrient, and Average Daily Gain of Local Indonesian Beef Cattle](#). *World Vet. J.*, 12 (4):

358-362. DOI: <https://dx.doi.org/10.54203/scil.2022.wvj45> Jentsch **8** W, Schweigel M, Weissbach F, Scholze H, Pitroff W, and Derno M (2007). Methane production in cattle calculated by the nutrient composition of the diet. *Archives of Animal Nutrition*, 61(1):

10-19. DOI: <https://www.doi.org/10.1080/17450390601106580> **7** [Jolazadeh AR,](#)

Dehghan-banadaky M, and Rezayazdi K (2015). Effects of soybean meal treated with tannins extracted from pistachio hulls on performance, ruminal fermentation, blood metabolites and nutrient digestion of Holstein bulls. *Animal Feed Science and Technology*, 203: 33-40. DOI: <https://www.doi.org/10.1016/j.anifeedsci.2015.02.005> Ku-Vera JC, Jiménez-Ocampo R, Valencia-Salazar SS, <sup>2</sup> Montoya-Flores MD, Molina-Botero IC, Arango J, Gómez-Bravo CA, Aguilar-Pérez CF, and Solorio-Sánchez FJ (2020). <sup>15</sup> Role of secondary plant metabolites on enteric methane mitigation in ruminants. *Frontiers in Veterinary Science*, 7: 584. DOI: <https://www.doi.org/10.3389/fvets.2020.00584> Lardy G (2018). Quality forage series: Forage nutrition for ruminants. North Dakota state university, 1250: 1-16. Available at: <https://www.ndsu.edu/agriculture/sites/default/files/2022-07/as1250.pdf> <sup>3</sup> Sampaio CB, Detmann E, Paulino MF, Filho SCV, de Souza MA, Lazzarini I, Rodrigues Paulino PV, and Oliveira FA (2010). Intake and digestibility in cattle fed low-quality tropical forage and supplemented with nitrogenous compounds. *Tropical Animal Health and Production*, 42: 1471-1479. Available at: <https://link.springer.com/article/10.1007/s11250-010-9581-7> Le Bourvellec C and Renard CMGC (2018). Interactions between polyphenols and macromolecules: Effect of tannin structure. *Encyclopedia of food chemistry*, pp. 515-521. DOI: <https://www.doi.org/10.1016/B978-0-08-100596-5.21486-8> Liu H, Puchala R, Leshure S, Gipson TA, Flythe MD, and Goetsch AL (2019). <sup>5</sup> Effects of lespedeza condensed tannins alone or with monensin, soybean oil, and coconut oil on feed intake, growth, digestion, ruminal methane emission, and heat energy by yearling Alpine doelings. *Journal of Animal Science*, 97(2): 885-899. DOI: <https://www.doi.org/10.1093/jas/sky452> Medion (2022). Wormectin Injeksi. Available at: <https://www.medionfarma.co.id/product/wormectin-injeksi/> Ningrat RWS, Zain M, Erpomen, and Suryani H (2018). <sup>6</sup> Effects of supplementation of different sources of tannins on nutrient digestibility, methane production and daily weight gain of beef cattle fed on ammoniated oil palm frond based diet. *International Journal of Zoological Research*, 14(1): 8-13. DOI: <https://www.doi.org/10.3923/ijzr.2018.8.13> Ramaiyulis (2021). Rumen un-

degraded dietary protein and TCA soluble protein with gambier leaf residue supplementation as a source of tannins in cattle feed supplement. **1 IOP Conference**

**Series: Earth and Environmental Science**, 759: 012045. DOI:

<https://www.doi.org/10.1088/1755-1315/759/1/012045> Ramaiyulis, Yulia E, Sari DK, and Nilawati (2021). **Effect of addition cattle feed supplement on in vitro fermentation, synthesis of microbial biomass, and methane production of rice straw fermentation basal diets. IOP Conference Series: Earth and Environmental Science**, 888: 012070. DOI:

<https://www.doi.org/10.1088/1755-1315/888/1/012070> Ramaiyulis, Ningrat RWS, Zain M, and Warly L (2019). **Optimization of rumen microbial protein synthesis by addition of gambier leaf residue to cattle feed supplement. Pakistan Journal of Nutrition**, 18(1): 12-19.

DOI: <https://www.doi.org/10.3923/pjn.2019.12.19> **12** Rira M, Morgavi DP, Popova M, Maxin G, and Doreau M (2022). **Microbial colonisation of tannin-rich tropical plants : Interplay between degradability , methane production and tannin disappearance in the rumen. Animal**, 16(8): 100589. DOI: <https://www.doi.org/10.1016/j.animal.2022.100589>

Rufino LMA, Detmann E, Gomes DÍ, dos Reis WLS, Batista ED, Filho SCV, and Paulino MF (2016). Intake, digestibility and nitrogen utilization in **19** **cattle fed tropical forage and supplemented with protein in the rumen, abomasum, or both. Journal of Animal Science and Biotechnology**, 7: 11. DOI: <https://www.doi.org/10.1186/s40104-016-0069-9> **3** Souza

**MA, Detmann E, Paulino MF, Sampaio CB, Lazzarini Í, and Filho SCV (2010). Intake, digestibility and rumen dynamics of neutral detergent fibre in cattle fed low-quality tropical forage and supplemented with nitrogen and/or starch. Tropical Animal Health and Production**, 42(6), 1299-1310. DOI: <https://www.doi.org/10.1007/s11250-010-9566-6>

Souza RA, Tempelman RJ, Allen MS, Weiss WP, Bernard JK, and VandeHaar MJ (2017). Predicting nutrient digestibility in highproducing dairy cows. *Journal of Dairy Science*, 101(2): 1123-1135. DOI: <https://www.doi.org/10.3168/jds.2017-13344> Stewart EK,

Beauchemin KA, Dai X, MacAdam JW, Christensen RG, and Villalba JJ (2019). Effect of tannin-containing hays on enteric methane emissions and nitrogen partitioning in beef cattle. **2** **Journal of Animal Science**, 97(8): 3286-3299. DOI:

<https://www.doi.org/10.1093/jas/skz206> van Kuijk S, Swiegers P, and Han Y (2022).

Hydroxychloride trace minerals improve apparent total tract nutrient digestibility in

Bonsmara beef cattle. *Livestock Science*, 256: 104820.

DOI: <https://www.doi.org/10.1016/j.livsci.2021.104820> Van Soest PJ, Robertson JB, and

Lewis BA (1991). Methods for dietary fiber, neutral detergent fiber, and nonstarch

polysaccharides in relation to animal nutrition. *Journal of Dairy Science*, 74(10):

3583-3597. DOI: [https://www.doi.org/10.3168/jds.S00220302\(91\)78551-2](https://www.doi.org/10.3168/jds.S00220302(91)78551-2) Wanapat M and

Kang S (2015). Cassava chip (*Manihot esculenta* Crantz) as an energy source for ruminant

feeding. *Animal Nutrition*, 1(4): 266-270. DOI:

<https://www.doi.org/10.1016/j.aninu.2015.12.001> 13 Watts N, Amann M, Arnell N, Ayeb-

Karlsson S, Beagley J, Belesova K, Boykoff M, Byass P, Cai W, Campbell-Lendrum D et

al. (2021). 14 The 2020 report of The Lancet Countdown on health and climate change:

Responding to converging crises. *The Lancet*, 397(10269): 129-170. DOI:

[https://www.doi.org/10.1016/S0140-6736\(20\)32290-X](https://www.doi.org/10.1016/S0140-6736(20)32290-X)

## Sources

1	<a href="http://cholar.google.co.uk/citations?user=R_RuyoM">http://cholar.google.co.uk/citations?user=R_RuyoM</a> INTERNET 4%
2	<a href="http://cd.ub.edu/602/Entrenamiento_nacional_bibliografico_de_aplicaciones_tecnologicas">http://cd.ub.edu/602/Entrenamiento_nacional_bibliografico_de_aplicaciones_tecnologicas</a> INTERNET 2%
	<a href="http://link.springer.com/rct/10.1007/1120-010-81">http://link.springer.com/rct/10.1007/1120-010-81</a> INTERNET 2%
4	<a href="http://www.rchgt.net/publication/2600-Th_Nutrient_content_Dry_Matter_Dg_tblty_rgnc_Matter_Dg_tblty_Totl_Dg_tbl_Nutrient_nd_N_Ru_n_Producton_of_Thr_Knd_of_ttl_F_dng_Modl">http://www.rchgt.net/publication/2600-Th_Nutrient_content_Dry_Matter_Dg_tblty_rgnc_Matter_Dg_tblty_Totl_Dg_tbl_Nutrient_nd_N_Ru_n_Producton_of_Thr_Knd_of_ttl_F_dng_Modl</a> INTERNET 2%
	<a href="http://www.rchgt.net/publication/222468_Effct_of_lpdz_condn_dtnn_nlon_or_th_onn_noyb_nol_ndcoconut_ol_onfdntk_groth_dgton_runl_thn_on_ndht_nrgy_by_yrlng_lpn_doling">http://www.rchgt.net/publication/222468_Effct_of_lpdz_condn_dtnn_nlon_or_th_onn_noyb_nol_ndcoconut_ol_onfdntk_groth_dgton_runl_thn_on_ndht_nrgy_by_yrlng_lpn_doling</a> INTERNET 1%
6	<a href="http://www.rchgt.net/publication/14886_Effct_of_applnton_ofdffrnt_ourc_of_tnn_n_on_nutrnt_dg_tblty_thn_producton_nd_dly_ghg_n_of_b_fcttl_fdon_on_tdol_pl_frondb_ddt">http://www.rchgt.net/publication/14886_Effct_of_applnton_ofdffrnt_ourc_of_tnn_n_on_nutrnt_dg_tblty_thn_producton_nd_dly_ghg_n_of_b_fcttl_fdon_on_tdol_pl_frondb_ddt</a> INTERNET 1%
	<a href="http://www.rchgt.net/publication/288_Influnc_of_Englnd_Tnn_n_ddton_on_Ru_n_Dgrdton_Kntc_of_Soy_Suc_Rdu">http://www.rchgt.net/publication/288_Influnc_of_Englnd_Tnn_n_ddton_on_Ru_n_Dgrdton_Kntc_of_Soy_Suc_Rdu</a> INTERNET 1%
8	<a href="http://cbrdg.org/cor/journal/dvnc-n-nl-bocnc/rct/b/gntc-control-of-grnhou-g-on/4DF64B6B00218DBE0102">http://cbrdg.org/cor/journal/dvnc-n-nl-bocnc/rct/b/gntc-control-of-grnhou-g-on/4DF64B6B00218DBE0102</a> INTERNET 1%
	<a href="http://cyberlink.ru/rct/n/ffct-of">http://cyberlink.ru/rct/n/ffct-of</a> INTERNET 1%
10	<a href="http://www.rchgt.net/publication/46_Idntfcion_of_ndophyte_nd_rhzo phr_bctr_n_rc_ryz_tv_n_th_xprnt_lfld_t_Py_kubuh_Stt_grcultura_Polychnc_Wt_Sustr_Ln don">http://www.rchgt.net/publication/46_Idntfcion_of_ndophyte_nd_rhzo phr_bctr_n_rc_ryz_tv_n_th_xprnt_lfld_t_Py_kubuh_Stt_grcultura_Polychnc_Wt_Sustr_Ln don</a> INTERNET 1%
11	<a href="http://www.rchgt.net/publication/414862_Th_Effct_of_Nutgv_Tnn_n_Myrtc_frgrn_outt_Protctng_gnt_on_Ln_Vtro_Nutrnt_Dg_tblty">http://www.rchgt.net/publication/414862_Th_Effct_of_Nutgv_Tnn_n_Myrtc_frgrn_outt_Protctng_gnt_on_Ln_Vtro_Nutrnt_Dg_tblty</a> INTERNET <1%
12	<a href="http://pub.dnbnl.nhgov/861/">http://pub.dnbnl.nhgov/861/</a> INTERNET <1%
1	<a href="http://crrp.org/frnc/frncpprpx?frncd412">http://crrp.org/frnc/frncpprpx?frncd412</a> INTERNET <1%

14	http://pub d ncb nl n hgov/ 2 8 / INTERNET <1%
1	http:// front r n org/ rt cl /10 8 /f uf 202166 00 /full INTERNET <1%
16	http:// r rchg t n t/publ c ton/24 44 6 _Ev lu ton_of_ n lyt c l_M thod _for_th _D t r n ton_ of_Mo tur _ rud _Prot n_ rud _F t_ nd_ rud _Fb r_n_D tll r _Dr d_Gr n _ th_Solubl INTERNET <1%
1	http:// lf lf ucd v du/ y po u /proc d ng /2014/14 S0 _Mc l n_ lf lf l prov nt.pdf INTERNET <1%
18	http:// po rn un v r ty c d/ c d c /c nt r-for-r rch- nd-co un ty- rv c / INTERNET <1%
1	http:// nt c chol r org/p p r/lnt k -d g t bl ty- nd-n trog n-ut lz ton-n-Ruf no- D t nn/ 6 c1d2 fb 06f 8 42 c 6 861c 8 8 b 2 INTERNET <1%
20	http:// p gov/cl t - nd c tor /cl t -ch ng - nd c tor - t o ph r c-conc ntr ton -gr nhou - g INTERNET <1%
21	http://cont ntc nc u du/nutr ton l-f d ng- n g nt-of- t-go t INTERNET <1%
22	http:// r rchg t n t/publ c ton/ 48 4081 _Soft r _for_D t _ n ly _n_SPSS_ n_ov r_v INTERNET <1%
2	http:// fl v n co /d/lg o/tr n l ton /Jl M rror/ ngl h/4 6_PP_ _2012_ ht l INTERNET <1%
24	http:// th -cryo ph r n t/pol c /co p t ng_ nt r t _pol cyht l INTERNET <1%
2	http://j pp ujour n l kb g/ rt cl _1882 0_0 c 2 bdbb f 0 1026224b6 pdf INTERNET <1%
26	http:// pn org/do/10 10 /pn 1 0814 110 INTERNET <1%

EXCLUDE CUSTOM MATCHES

OFF

EXCLUDE QUOTES OFF

EXCLUDE BIBLIOGRAPHY OFF